

ARTIGO

NEW ANTAGONIST AGENTS OF NEUROPEPTIDE Y RECEPTORS

Ignacio Aldana, Isabel Rivero, Argimiro Rivero, Patricio Huenchunir, Carmen Frigola, Maria Luisa Alonso e Antonio Monge

Department of Pharmaceutical Chemistry - CIFA - University of Navarra - Pamplona - Spain

D. H. Caignard and P. Renard

ADIR - Courbevoie Cedex - France

Recebido em 2/9/99; aceito em 20/4/00

In the CNS, NPY has been implicated in obesity and feeding, endocrine function and metabolism. Potent and selective rNPY antagonists will be able to probe the merits of this approach for the treatment of obesity. We report the synthesis and preliminary evaluation of some hydrazide derivatives as antagonists of rNPY.

Keywords: obesity; energy metabolism; weight reduction; neuropeptide Y.

INTRODUCTION

The Neuropeptide Y (NPY) is a peptide of 36 amino acids, similar to the peptide YY (PYY) and to the pancreatic polypeptides (PP)¹. At first, separated from the pig brain, the NPY is well-distributed in the mammals at the level of the central and peripheric nervous system. This neurotransmitter is present in strong concentrations in the nervous fiber of the brain, but equally as strong in the heart, the sympathetic glands, the blood vessels and the gastrointestinal tract. It is responsible for diverse physiological effects that exert themselves by way of specific receptors (Y). These Y receptors form a heterogeneous group and six subtypes have been identified up to now: Y1 to Y6². The NPY is involved in the control of appetite, strongly stimulating the intake of food, or performing a regulatory role on the HPA (hypothalamic-pituitary-adrenal) axis³. It also presents anxiolytic and sedative properties that induce an increase in the blood pressure and affects the circadian rhythm⁴.

The search for agonists and antagonists of NPY could yield compounds which could be used in the treatment of pathologies related to food behavior or energy equilibrium disorders, such as diabetes, obesity, bulimia and anorexia. Also in the treatment of arterial hypertension, anxiety, depressions, epilepsy and sleeping disorders.

The orexigenic effects of NPY could be mediated by its receptor NPY1 subtype⁵, which makes it possible to consider that is involved in the stimulating effect of the appetite of the neuropeptide Y. Recently, ligands of this receptor have been described. The first compound of nonpeptide structure with potent antagonist activity of the receptor NPY1, is BIBP 3226 (Fig. 1), which was designed using the region C-terminal of the NPY⁶. Recent studies appear to confirm an important implication of the receptor NPY5 subtype in the control of food ingestion⁷. The first Y5-selective antagonists appeared in the patent literature in mid-1997⁸. These compounds have something in common: an amino acid, phenylalanine (SR 120819A) and arginine (BIBP 3226) respectively, in the center part of the molecule. Also, the amino and carboxyl groups from the amino acid remain functionalized in the form of amide. Likewise, the first selective NPY5 antagonists are appearing. One such example is JCF 104 (2-(naphthalen-1-yl)-3-phenylpropane-1,2-diamine)⁹ (Fig. 1), which carries a rest of 2-amino-3-phenylpropanamine, closely related to the aforementioned systems. Our objective is to search for new compounds that are non-peptidic selective antagonists of the Y5 receptor subtype. This line of research implies the presence of a phenylalanine and/or methyltyrosine rest in the central part of a molecule in

which the carboxyl and amine groups of the amino acid are functionalized as amide and carbamate respectively (Fig. 2).




Figure 1. Antagonists of the NPY receptors.




Figure 2. General structure for the synthesized compounds.

In this work, the first experimental results are offered. All the synthesized compounds were evaluated for their antagonist action of NPY receptors. Their action was evaluated on the NPY Y1, Y2, Y5 subtype receptors.

CHEMISTRY

Chemicals, methods and apparatus

General laboratory chemicals were purchased from Merck, Sigma, Janssen, and Scharlau. L-phenylalanine, L-methyltyrosine, HOBt and EDC was purchased from Fluka.

All the new compounds were characterized by elemental analysis, IR and ^1H NMR. The IR spectra are recorded on a Perkin-Elmer FT 681 using KBr pellets. The ^1H NMR spectra are obtained on a Bruker Model AC-200E (200 MHz) with Me_4Si as the internal standard, and at a concentration of approximately 0.1 g/mL. The mass spectra were obtained on a Hewlett-Packard HP-5890 (GC/HPLC/DIP) instrument. All spectra were consistent with assigned structures. Thin layer chromatography was performed on aluminum sheets precoated with silica gel (HF 254, Merck). The developed chromatograms were viewed under UV light or iodine revelation. Melting points were determined on a Mettler FP82 hot stage apparatus equipped with a FP800/FP80 processor and an Olympus 8091 microscope provided with a video system and were uncorrected. Elemental analyses of vacuum-dried samples were obtained on a Carlo Erba 1106 elemental analyzer (over P_2O_5 at 1-2 mmHg, 24 h at 60-80°C). Results are within 0.4% of theoretical values unless otherwise indicated.

Synthesis of the compounds

General procedure for the preparation of 3-H and 3-(4-methoxyphenyl)propionic acid derivatives. a-b

The appropriate amino acid was dissolved in NaOH 4 M. The solution was cooled to 0°C in an ice bath. A solution of the benzyl chloroformate (equimolar amount) in dry CH_2Cl_2 and NaOH 4 M were added alternately and portionwise, with stirring and cooling, during 30 minutes. The mixture was allowed to warm with intermittent shaking for 1 h. The mixture was extracted with ethyl ether (2-15 mL). The aqueous layer was acidified with diluted HCl (1:1) to pH 2. The solid was filtered and washed with ethyl ether (3x10 mL) in order to obtain the product as a solid.

(S)-2-Benzylloxycarbonylamino-3-phenylpropanoic acid (a)

From L-Phe (13.20 g, 80.00 mmol) and benzyl chloroformate (11.25 g, 80.00 mmol). A white solid (20.87 g, 87%) mp 84.9-85.4: IR (KBr, cm^{-1}) 3329 (s, N-H), 3033 (m, aromatic C-H), 2942 (m, aliphatic C-H), 1696 (vs, carbamate and carboxylic acid C=O), 748 and 697 (m, aromatic monosubstitution); ^1H -NMR (CDCl_3 , δ) 2.74-3.10 (d, d, d, d, 2H, $\text{CH}_2\text{-CH}$, J=5.5, 5.3, 6.2, 6.4 Hz), 4.06-4.17 (dd, 1H, CH , J=6.0/13.8 Hz), 4.93 (s, 2H, $\text{CH}_2\text{-O}$), 6.42 (d, 1H, NH , J=7.7 Hz), 7.05-7.35 (m, 10H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-CH}$ and $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$), 10.18 (ws, 1H, OH); MS m/z 107.05 (10.56), 91.05 (100), 76.95 (12.00); Anal. ($\text{C}_{17}\text{H}_{17}\text{NO}_4$) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-(4-methoxyphenyl)propanoic acid (b)

From O-Methyl-L-Tyr (0.50 g, 2.50 mmol) and benzyl chloroformate (0.36 mL, 2.56 mmol). A white solid (0.65 g, 77%) mp 111.1-113.4: IR (KBr, cm^{-1}) 3305 (s, N-H), 3033 (m, aromatic C-H), 2956 (m, aliphatic C-H), 1696 (vs, carbamate and carboxylic acid C=O), 746 and 695 (aromatic monosubstitution); ^1H -NMR (CDCl_3 , δ) 3.01-3.22 (d, t, d, 2H, $\text{CH}_2\text{-CH}$, J=5.9, 6.0, 5.5 Hz), 3.78 (s, 3H, OCH_3), 4.62-4.71 (dd, 1H, CH , J=5.9/13.5 Hz), 5.11 (s, 2H, $\text{CH}_2\text{-O}$), 5.21 (d, 1H, NH , J=8.1 Hz), 6.82 (d, 2H, H_2 and H_6 of $\text{C}_6\text{H}_4\text{-OCH}_3$, J=8.5 Hz), 7.07 (d, 2H, H_3 and H_5 of $\text{C}_6\text{H}_4\text{-OCH}_3$, J=8.5 Hz), 7.35 (s, 5H, $\text{C}_6\text{H}_5\text{-CH}_2$); MS m/z 329.05 (1.50, M^+), 121.15 (100), 91.05 (12.09), 77.05 (15.22); Anal. ($\text{C}_{18}\text{H}_{19}\text{NO}_5$) C, H, N.

General procedure for the preparation of final compounds (1-13)

A mixture of the appropriate propanoic acid derivative and HOBT in dry CH_2Cl_2 at 0°C, under N_2 , was treated with EDC. After 1h at 0°C, a solution of the appropriate hydrazide or hydrazine, in dry CH_2Cl_2 was added (equimolar amount of HOBT, EDC, and hydrazide or hydrazine was used). The reaction was stirred at

0°C for 1 h and then at room temperature for 24 h. The solvent was evaporated. The residue was redissolved in ethyl acetate or CH_2Cl_2 and washed with water (3x20 mL). The organic phase was dried over Na_2SO_4 and evaporated under reduced pressure. The obtained residue was washed with ethyl ether in order to obtain the product as a solid.

(S)-2-Benzylloxycarbonylamino-3-phenyl-N'-phenylpropionohydrazide (1)

From (a) (1.00 g, 3.34 mmol), HOBT (0.58 g, 3.81 mmol), EDC (0.73 g, 3.81 mmol) and phenylhydrazine (0.37 mL, 3.81 mmol). A beige solid (0.53 g, 38%) mp 175.0-177.1: IR (KBr, cm^{-1}) 3297 and 3244 (m, N-H), 3032 (w, aromatic C-H), 2920 (w, aliphatic C-H), 1688 (s, carbamate C=O), 1656 (s, hydrazide C=O), 746 and 696 (m, aromatic monosubstitution); ^1H -NMR (CDCl_3 , δ) 3.07 (d, 2H, $\text{CH}_2\text{-CH}$, J=7.3 Hz), 4.45-4.60 (dd, 1H, CH , J=7.4/15.1 Hz), 5.06 (s, 2H, $\text{Ph-CH}_2\text{-O}$), 5.49 (d, 1H, NH-CH , J=8.5 Hz), 5.97 (d, 1H, NH-NH-Ph , J=3.6 Hz), 6.50 (d, 2H, H_2 and H_6 of $\text{C}_6\text{H}_5\text{-NH}$, J=7.9 Hz), 6.85 (t, 1H, H_4 of $\text{C}_6\text{H}_5\text{-NH}$, J=7.3 Hz), 7.07-7.35 (m, 12H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$, $\text{C}_6\text{H}_5\text{-CH}_2\text{-CH}$ and H_3 and H_5 of $\text{C}_6\text{H}_5\text{-NH}$), 8.01 (ws, 1H, CO-NH-NH); MS m/z 389.20 (2.90, M^+), 107.15 (51.51), 91.05 (100); Anal. ($\text{C}_{23}\text{H}_{23}\text{N}_3\text{O}_3$) C, H, N.

(S)-N'-Benzoyl-2-benzylloxycarbonylamino-3-phenylpropionohydrazide (2)

From (a) (1.00 g, 3.34 mmol), HOBT (0.56 g, 3.68 mmol), EDC (0.71 g, 3.68 mmol) and phenylhydrazide (0.50 g, 3.68 mmol). A beige solid (0.20 g, 14%) mp 172.2-174.5: IR (KBr, cm^{-1}) 3285 and 3205 (m, N-H), 3030 (w, aromatic C-H), 1699 (s, carbamate C=O), 1610 (s, hydrazide C=O), 744 and 696 (m, aromatic monosubstitution); ^1H -NMR (CDCl_3 , δ) 2.97-3.24 (d, d, d, d, 2H, $\text{CH}_2\text{-CH}$, J=8.3, 8.2, 5.9, 6.1 Hz), 4.87-5.06 (m, 3H, CH and $\text{CH}_2\text{-O}$), 5.98 (ws, 1H, NH-CH), 7.09-7.38 (m, 12H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$, $\text{C}_6\text{H}_5\text{-CH}_2\text{-CH}$ and H_3 and H_5 of $\text{C}_6\text{H}_5\text{-CO}$), 7.48 (t, 1H, H_4 of $\text{C}_6\text{H}_5\text{-CO}$, J=7.2 Hz), 7.78 (d, 2H, H_2 and H_6 of $\text{C}_6\text{H}_5\text{-CO}$, J=7.8 Hz), 9.52 (ws, 1H, CH-CO-NH-NH-CO), 9.96 (ws, 1H, CO-NH-NH-CO-Ph); MS m/z 282.15 (14.37), 254.15 (3.91), 105.00 (54.12), 91.05 (100); Anal. ($\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_4$) C, H, N.

(S)-2-Benzylloxycarbonylamino-N'-(2-indolylcarbonyl)-3-phenylpropionohydrazide (3)

From (a) (0.20 g, 0.67 mmol), HOBT (0.12 g, 0.76 mmol), EDC (0.14 g, 0.76 mmol) and indol-2-hydrazide (0.14 g, 0.76 mmol). A beige solid (0.15 g, 49%) mp 176.3-177.5: IR (KBr, cm^{-1}) 3412 (vs, N-H), 3031 (w, aromatic C-H), 2953 (w, aliphatic C-H), 1697 (vs, carbamate C=O), 1666 (vs, hydrazide C=O), 745 and 697 (vs, aromatic monosubstitution); ^1H -NMR (DMSO-d_6 , δ) 2.90 (t, 1H, $\text{CH}_A\text{H}_B\text{-CH}$, J=11.2 Hz), 3.20 (d, 1H, $\text{CH}_A\text{H}_B\text{-CH}$, J=11.8 Hz), 4.30-4.60 (m, 1H, CH), 4.98 (s, 2H, $\text{CH}_2\text{-O}$), 6.95-7.55 (m, 13H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$, $\text{C}_6\text{H}_5\text{-CH}_2\text{-CH}$, H_3 , H_5 and H_6 of indol-CO), 7.55-7.80 (m, 2H, H_4 and H_7 of indol-CO), 10.36 (s, 1H, CH-CO-NH-NH-CO), 10.56 (s, 1H, CO-NH-NH-CO-Ph), 11.77 (s, 1H, NH of indole); MS m/z 348.00 (5.41), 143.95 (42.20), 90.95(100); Anal. ($\text{C}_{26}\text{H}_{24}\text{N}_4\text{O}_4$) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-phenyl-N'-(3-pyridylcarbonyl)propionohydrazide (4)

A mixture of (a) (0.50 g, 1.67 mmol), HOBT (0.28 g, 1.84 mmol), EDC (0.35 g, 1.84 mmol) and pyridyl-3-hydrazide (0.25 g, 1.84 mmol). A white solid (0.16 g, 22%) mp 174.0-176.0: IR (KBr, cm^{-1}) 3419 (s, OH), 3284 (s, NH), 3028 (w, aromatic C-H), 1691 (s, carbamate C=O), 1659 (s, hydrazide C=O), 745 and 697 (s, aromatic monosubstitution); ^1H -NMR (DMSO-d_6 , δ) 2.84 (t, 1H, $\text{CH}_A\text{H}_B\text{-CH}$, J=12.3 Hz), 3.13 (d, 1H, $\text{CH}_A\text{H}_B\text{-CH}$, J=11.9 Hz), 4.31-4.52 (m, 1H, CH), 4.95 (s, 2H, $\text{CH}_2\text{-O}$), 7.12-7.44 (m, 10H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-CH}$ and $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$), 7.53-7.59 (m, 1H, H_5 of pyridyl-CO), 7.71 (d, 1H, H_6 of pyridyl-CO, J=8.6

Hz), 8.25 (d, 1H, NH-CH, J=7.9 Hz), 8.77 (d, 1H, **H**₄ of pyridyl-CO, J=3.8 Hz), 9.06 (s, 1H, **H**₂ of pyridyl-CO), 10.39 (s, 1H, CH-CO-NH-NH-CO), 10.72 (s, 1H, CO-NH-NH-CO-pyridine); MS *m/z* 418.05 (7.39, M⁺), 254.15 (9.12), 91.00 (100); Anal. (C₂₃H₂₂N₄O₄) C, H, N.

(S)-2-Benzylloxycarbonylamino-N'-(3-indolylacethyl)-3-phenylpropionohydrazide (5)

From **(a)** (0.40 g, 1.34 mmol), HOBT (0.23 g, 1.47 mmol), EDC (0.28 g, 1.47 mmol) and 3-indolylacetylhydrazide (0.28 g, 1.47 mmol). A white solid (0.39 g, 57%) mp 156.7-158.5: IR (KBr, cm⁻¹) 3279 and 3230 (vs, N-H), 3055 (m, aromatic C-H), 2920 (w, aliphatic C-H), 1697 (s, carbamate C=O), 1667 and 1608 (vs, hydrazide C=O), 747 and 697 (s, aromatic monosubstitution); ¹H-NMR (DMSO-d₆, δ) 2.80 (t, 1H, CH_AH_B-CH, J=12.2 Hz), 3.05 (d, 1H, CH_AH_B-CH, J=12.5 Hz), 3.93 (s, 2H, CO-CH₂-indole), 4.31-4.52 (m, 1H, CH), 4.95 (s, 2H, CH₂-O), 6.95-7.20 (m, 3H, **H**₂, **H**₅ and **H**₆ of indolyl-CH₂), 7.20-7.45 (m, 10H, C₆H₅-CH₂-CH and C₆H₅-CH₂-O), 7.50-7.70 (m, 2H, **H**₄ and **H**₇ of indolyl-CH₂), 10.41 (s, 2H, CO-NH-NH-CO), 11.17 (s, 1H, NH of indolyl-CH₂); MS *m/z* 392.05 (11.81), 157.15 (86.51), 130.15 (100), 108.15 (48.18); Anal. (C₂₇H₂₆N₄O₄) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-phenyl-N-(3-pyrazolyl)propionamide (6)

From **(a)** (0.50 g, 1.67 mmol), HOBT (0.28 g, 1.84 mmol), EDC (0.35 g, 1.84 mmol) and 3-aminopyrazole (0.15 g, 1.84 mmol). A white solid (0.23 g, 34%) mp 113.3-114.9: IR (KBr, cm⁻¹) 3268 (vs, N-H), 3032 (s, aromatic C-H), 2953 (s, aliphatic C-H), 1713 (vs, carbamate C=O), 1674 (vs, amide C=O), 738 and 697 (s, aromatic monosubstitution); ¹H-NMR (CDCl₃, δ) 3.05-3.28 (m, 2H, CH₂-CH), 4.79-4.93 (m, 1H, CH), 5.06 (dd, 2H, CH₂-O, J=12.4/28.0 Hz), 6.61 (d, 1H, NH-CH, J=9.5 Hz), 6.84 (s, 1H, **H**₄ of pyrazolyl-NH), 7.15-7.36 (m, 11H, **H**₅ of pyrazolyl-NH, C₆H₅-CH₂-CH and C₆H₅-CH₂-O), 11.53 (ws, 1H, CH-CO-NH), 11.72 (ws, 1H, NH of pyrazolyl-NH); MS *m/z* 364.05 (0.75, M⁺), 256.15 (51.31), 108.00 (26.49), 91.00 (100); Anal. (C₂₀H₂₀N₄O₃·1H₂O) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-(4-methoxyphenyl)-N'-phenylpropionohydrazide (7)

From **(b)** (0.65 g, 1.98 mmol), HOBT (0.33 g, 2.18 mmol), EDC (0.42 g, 2.18 mmol) and phenylhydrazine (0.20 mL, 2.18 mmol). A beige solid (0.62 g, 75%) mp 183.0-185.0: IR (KBr, cm⁻¹) 3299 and 3241 (m, N-H), 3038 (w, aromatic C-H), 2963 (w, aliphatic C-H), 1688 (s, carbamate C=O), 1657 (s, amide C=O), 739 and 694 (m, aromatic monosubstitution); ¹H-NMR (CDCl₃, δ) 2.95-3.10 (m, 2H, CH₂-CH), 3.77 (s, 3H, OCH₃), 4.46-4.50 (m, 1H, CH), 5.06 (s, 2H, CH₂-O), 5.45 (d, 1H, NH-CH, J=7.5 Hz), 5.98 (d, 1H, NH-NH-Ph, J=3.2 Hz), 6.51 (d, 2H, **H**₂ and **H**₆ of C₆H₅-NH, J=7.8 Hz), 6.77-6.88 (m, 3H, **H**₄ of C₆H₅-NH and **H**₂ and **H**₆ of C₆H₄-OCH₃), 6.99-7.11 (m, 4H, **H**₃ and **H**₅ of C₆H₅-NH and **H**₃ and **H**₅ of C₆H₄-OCH₃), 7.25 (s, 5H, C₆H₅-CH₂), 7.94 (ws, CH-CO-NH-NH); MS *m/z* 419.20 (11.70, M⁺), 150.15 (29.57), 121.15 (100), 91.05 (74.84); Anal. (C₂₄H₂₅N₃O₄) C, H, N.

(S)-N'-Benzoyl-2-benzylloxycarbonylamino-3-(4-methoxyphenyl)propionohydrazide (8)

From **(b)** (0.65 g, 1.98 mmol), HOBT (0.33 g, 2.18 mmol), EDC (0.42 g, 2.18 mmol) and phenylhydrazide (0.30 g, 2.18 mmol). A beige solid (0.36 g, 41%) mp 208.0-209.2: IR (KBr, cm⁻¹) 3280 (s, N-H), 3033 (w, aromatic C-H), 2925 (w, aliphatic C-H), 1689 (vs, carbamate C=O), 1655 (vs, hydrazide C=O), 824 (m, aromatic *p*-disubstitution), 745 and 695 (m, aromatic monosubstitution); ¹H-NMR (CDCl₃, δ) 2.90-3.12 (m, 2H, CH₂-CH), 3.68 (s, 3H, OCH₃), 4.50-4.60 (m, 1H, CH), 4.91-5.06 (dd, 2H, CH₂-O, J=12.3/18.8 Hz), 5.53 (ws, 1H, NH-CH), 6.71 (d, 2H,

H₃ and **H**₅ of C₆H₄-OCH₃, J=8.5 Hz), 7.04 (d, 2H, **H**₂ and **H**₆ de C₆H₄-OCH₃, J=8.4 Hz), 7.10-7.25 (m, 5H, C₆H₅-CH₂-O), 7.34 (t, 2H, **H**₃ and **H**₅ of C₆H₅-CO, J=7.1 Hz), 7.46 (t, 1H, **H**₄ of C₆H₅-CO, J=7.3 Hz), 7.73 (d, 2H, **H**₂ and **H**₆ of C₆H₅-CO, J=7.5 Hz), 8.90-9.30 (ws, 2H, CO-NH-NH-CO); MS *m/z* 339.05 (4.01), 150.15 (1.69), 121.15 (100), 115.00 (40.58), 91.05 (63.37); Anal. (C₂₅H₂₅N₃O₃) C, H, N.

(S)-2-Benzylloxycarbonylamino-N'-(2-indolylcarbonyl)-3-(4-methoxyphenyl)propionohydrazide (9)

From **(b)** (0.30 g, 0.91 mmol), HOBT (0.16 g, 1.04 mmol), EDC (0.20 g, 1.04 mmol) and indol-2-hydrazide (0.19 g, 1.04 mmol). A beige solid (0.20 g, 48%) mp 185.5-186.5: IR (KBr, cm⁻¹) 3359 (vs, N-H), 3038 (w, aromatic C-H), 2963 (w, aliphatic C-H), 1707 (vs, carbamate C=O), 1654 and 1606 (vs, hydrazide C=O), 822 (s, aromatic *p*-disubstitution), 743 (vs, aromatic monosubstitution); ¹H-NMR (DMSO-d₆, δ) 2.81 (t, 1H, CH_AH_B-CH, J=12.2 Hz), 3.11 (d, 1H, CH_AH_B-CH, J=13.7 Hz), 3.74 (s, 3H, OCH₃), 4.38 (ws, 1H, CH), 4.97 (s, 2H, CH₂-O), 6.87 (d, 2H, **H**₃ and **H**₅ of C₆H₄-OCH₃, J=8.2 Hz), 7.07 (t, 1H, **H**₅ of indol-CO, J=7.2 Hz), 7.10-7.40 (m, 9H, C₆H₅-CH₂, **H**₂ and **H**₆ of C₆H₄-OCH₃ and **H**₆ and **H**₃ of indol-CO), 7.47 (d, 1H, **H**₄ of indol-CO, J=8.0 Hz), 7.66 (d, 1H, **H**₇ of indol-CO, J=8.4 Hz), 7.98 (d, 1H, NH-CH, J=8.3 Hz), 10.31 (s, 1H, CH-CO-NH-NH-CO), 10.52 (s, 1H, CO-NH-NH-CO-indole), 11.74 (s, 1H, NH of indol-CO); MS *m/z* 335.15 (7.99), 144.05 (55.08), 121.05 (100); Anal. (C₂₇H₂₆N₄O₅) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-(4-methoxyphenyl)-N'-(3-pyridylcarbonyl)propionohydrazide (10)

From **(b)** (0.30 g, 0.91 mmol), HOBT (0.15 g, 0.99 mmol), EDC (0.19 g, 0.99 mmol) and pyridyl-3-hydrazide (0.14 g, 0.99 mmol). A white solid (0.25 g, 61%) mp 184.0-186.0: IR (KBr, cm⁻¹) 3279 (s, N-H), 3033 (w, aromatic C-H), 2958 (w, aliphatic C-H), 1691 (s, carbamate C=O), 1660 (s, hydrazide C=O), 813 (w, aromatic *p*-disubstitution), 751 and 704 (m, aromatic monosubstitution); ¹H-NMR (CDCl₃, δ) 2.93-3.16 (m, 2H, CH₂-CH), 3.72 (s, 3H, OCH₃), 4.55-4.69 (m, 1H, CH), 4.95-5.09 (dd, 2H, CH₂-O, J=12.3/17.3 Hz), 5.66 (ws, 1H, NH-CH), 6.76 (d, 2H, **H**₃ and **H**₅ of C₆H₄-OCH₃, J=8.4 Hz), 7.08 (d, 2H, **H**₂ and **H**₆ of C₆H₄-OCH₃, J=8.2 Hz), 7.20-7.36 (m, 6H, C₆H₅-CH₂ and **H**₅ of pyridyl-CO), 8.07 (d, 1H, **H**₆ of pyridyl-CO, J=7.9 Hz), 8.70 (d, 1H, **H**₄ of pyridyl-CO, J=4.8 Hz), 9.01 (s, 1H, **H**₂ of pyridyl-CO), 9.37 (ws, 2H, CH-CO-NH-NH-CO-pyridine); MS *m/z* 297.15 (6.82), 121.05 (100), 91.15 (49.17); Anal. (C₂₄H₂₄N₄O₅) C, H, N.

(S)-2-Benzylloxycarbonylamino-3-(4-methoxyphenyl)-N'-phenylsulphonylpropionohydrazide (11)

From **(b)** (0.30 g, 0.91 mmol), HOBT (0.15 g, 0.99 mmol), EDC (0.19 g, 0.99 mmol) and benzenesulfonyl hydrazide (0.17 g, 0.99 mmol). A beige solid (0.17 g, 38%) mp 167.0-169.0: IR (KBr, cm⁻¹) 3303 (s, N-H), 3063 (w, aromatic C-H), 2962 (w, aliphatic C-H), 1691 (s, carbamate C=O), 1653 (s, hydrazide C=O), 1360 and 1173 (s, SO₂), 834 (m, aromatic *p*-disubstitution), 734 and 689 (m, aromatic monosubstitution); ¹H-NMR (CDCl₃, δ) 2.56-2.70 (m, 2H, CH₂-CH), 3.69 (s, 3H, OCH₃), 4.19-4.27 (m, 1H, CH), 5.00 (s, 2H, CH₂-O), 6.69 (d, 2H, **H**₃ and **H**₅ of C₆H₄-OCH₃, J=8.5 Hz), 6.86 (d, 2H, **H**₂ and **H**₆ of C₆H₄-OCH₃, J=8.5 Hz), 7.19-7.53 (m, 9H, C₆H₅-CH₂ and **H**₂, **H**₃, **H**₅ and **H**₆ of C₆H₅-SO₂), 7.79 (d, 1H, **H**₄ of C₆H₅-SO₂, J=7.4 Hz), 9.05 (ws, 2H, CO-NH-NH-SO₂); MS *m/z* 332.15 (7.19), 284.20 (1.36), 121.05 (100), 91.15 (94.19); Anal. (C₂₄H₂₅N₃SO₆) C, H, N.

(S)-2-Benzylloxycarbonylamino-N'-(3-indolylacethyl)-3-(4-methoxyphenyl)propionohydrazide (12)

From **(b)** (0.57 g, 1.76 mmol), HOBT (0.30 g, 1.94 mmol),

EDC (0.37 g, 1.94 mmol) and 3-indolylacetohydrazide (0.37 g, 1.94 mmol). A beige solid (0.51 g, 53%) mp 194.0-195.5: IR (KBr, cm^{-1}) 3410 and 3272 (vs, N-H), 3034 (s, aromatic C-H), 2949 (m, aliphatic C-H), 1703 (vs, carbamate C=O), 1602 (vs, hydrazide C=O), 824 (s, aromatic *p*-disubstitution), 735 and 698 (vs, aromatic monosubstitution); $^1\text{H-NMR}$ (DMSO- d_6 , δ) 2.73 (t, 1H, $\text{CH}_A\text{H}_B\text{-CH}$, $J=12.0$ Hz), 2.98 (d, 1H $\text{CH}_A\text{H}_B\text{-CH}$, $J=12.3$ Hz), 3.61 (s, 2H, $\text{CH}_2\text{-indole}$), 3.71 (s, 3H, OCH_3), 4.30 (ws, 1H, CH), 4.95 (s, 2H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$), 6.84 (d, 2H, H_3 and H_5 of $\text{C}_6\text{H}_4\text{-OCH}_3$, $J=7.3$ Hz), 6.90-7.40 (ws, 11H, $\text{C}_6\text{H}_5\text{-CH}_2\text{-O}$, H_2 and H_6 of $\text{C}_6\text{H}_4\text{-OCH}_3$, H_2 , H_5 and H_6 of indolyl- CH_2 and NH-CH), 7.55-7.75 (m, 2H, H_4 and H_7 of indolyl- CH_2), 10.17 (s, 2H, CO-NH-NH-CO), 10.92 (s, 1H, NH of indolyl- CH_2); MS m/z 500.20 (2.42, M^+), 392.05 (4.80), 157.15 (100), 130.15 (77.01), 108.15 (68.11); Anal. ($\text{C}_{28}\text{H}_{28}\text{N}_4\text{O}_5$) C, H, N.

(*S*)-2-Benzoyloxycarbonylamino-3-(4-methoxyphenyl)-*N*-(3-pyrazolyl)propionamide (**13**)

From (**b**) (0.50 g, 1.52 mmol), HOBT (0.26 g, 1.67 mmol), EDC (0.32 g, 1.67 mmol) and 3-aminopyrazole (0.14 g, 1.67 mmol). A white solid (0.28 g, 39%) mp 167.7-169.2: IR (KBr, cm^{-1}) 3268 (vs, N-H), 3032 (s, aromatic C-H), 2953 (s, aliphatic C-H), 1713 (vs, carbamate C=O), 1674 (vs, amide C=O), 813 (m, aromatic *p*-disubstitution), 738 and 697 (s, aromatic monosubstitution); $^1\text{H-NMR}$ (CDCl_3 , δ) 2.90-3.20 (m, 2H, $\text{CH}_2\text{-CH}$), 3.69 (s, 3H, OCH_3), 4.65-4.85 (m, 1H, CH), 4.90-5.20 (m, 2H, $\text{C}_6\text{H}_5\text{-CH}_2$), 6.63-6.81 (m, 4H, H_4 of pyrazolyl-NH, H_3 and H_5 of $\text{C}_6\text{H}_4\text{-OCH}_3$ and NH-CH), 7.06 (d, 2H, H_2 and H_6 of $\text{C}_6\text{H}_4\text{-OCH}_3$, $J=7.8$ Hz), 7.15-7.35 (m, 6H, H_5 of pyrazolyl-NH and $\text{C}_6\text{H}_5\text{-CH}_2$), 11.50 (ws, 1H, CH-CO-NH), 11.65 (ws, 1H, NH of pyrazolyl-NH); MS m/z 286.15 (16.41), 121.15 (100), 108.15 (49.21); Anal. ($\text{C}_{21}\text{H}_{22}\text{N}_4\text{O}_4$) C, H, N.

PHARMACOLOGY

Binding assays

Y1 binding assay

Binding of iodinated peptide YY (NEN) was carried out as described by Serradeil-Le Gal¹⁰ and modified as follows: incubations were performed at 30°C for 90 min. with various competitor concentrations in Buffer A (Herpes/NaOH 20 mM, MgSO_4 0.81 mM, KH_2PO_4 220 mM, CaCl_2 1.26 μM , MgSO_4 0.81 mM and bovine serum albumin 0.1%) with SK-N-MC cell membranes (50 μg of protein/mL of assay) in a total volume of 500 μL . Non-specific binding was determined in the presence of 1 μM NPY. The reaction was then stopped by filtration, the filters (GF/B, Whatman, pre-coated in 0.3% PEI) were extensively washed with buffer A, and counted in a gamma counter (Packard).

Y2 binding assay

Binding of iodinated peptide YY was conducted as described by Fuhlendorff *et al.*¹¹ on membrane prepared from KAN-TS cells.

Y5 binding assay


Binding of iodinated peptide YY (NEN) was carried out as described by Hu *et al.*¹². In brief, COS cells transfected with the human Y5 NPY receptor were lysed and the membranes prepared by differential centrifugation. These membranes contained about 2 pmol per mg of protein of this receptor. Incubations were performed in 500 μL comprising, 20 pM final of [^{125}I]PPYY in 50 μL . 400 μL of membrane suspension (0.15 mg/mL) and competitor

dilutions in 50 μL , at 30°C for 2 h. The reaction was stopped by filtration through GF/C filters (Whatman).

Note: Given the nature of the experiment the obtained values are within an acceptable variability range for this type of experiments, and therefore the standard errors are not included in the report.

RESULTS AND DISCUSSION


The carbamate derivatives were obtained as shown in Scheme 1. Formation of the carbamate is carried out by reaction of the benzyl chloroformate with the appropriate amino acid dissolved in a solution of NaOH 4 M. The use of strong basic mediums is necessary in order to solubilize the amino acids. In the reactions of the acid group of the amino acid fragment with amines, hydrazines or hydrazides, the work was carried out with *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC) and 1-hydroxybenzotriazole (HOBT), and with dichloromethane as the solvent of low dielectric constant. In each case, the corresponding amides or hydrazides were obtained. The reaction of the amino group of the amino acid does not lead to inversion of the configuration, but the reaction of the carboxyl group may give rise to different percentages of racemization. In order to avoid this, the reaction has been carried out with coupling agents that avoid racemization according to studies carried out by other authors¹³.



Scheme 1. Synthesis of hydrazide derivatives

The activity of compounds was evaluated by binding assay conducted as described by Félétou *et al.*¹⁴. Initially, the antagonist activity of the compounds was evaluated in a binding assay for Y1, Y2 and Y5 subtypes receptors. Competitive binding assays were performed using [^{125}I]-PPY as the radioligand, at concentrations which varied from 15 to 65 pM. The nonspecific fraction is measured in the presence of a concentration of 1 μM of NPY. The cells are incubated during a time which varied from 1 to 2 hours depending upon the classes, and the radioactivity is collected after filtration with a GF/C filter treated with PEI 0.1%, before being measured. The data corresponding to the products evaluated are shown in Table 1.

The results of affinity by the NPY subtype receptors Y1, Y2, Y5, from which the antagonist activity of the synthesized molecules has been determined, indicate that the molecules which were prepared from L-phenylalanine have better behavior than those prepared from L-methyltyrosine. The compound **5** ($\text{IC}_{50} = 4$ μM), which has a heterocyclic system of indole manifests an important level of activity. These initial results have convinced us to continue our work in the optimization of the starting base structure, searching for new highly selective nonpeptide antagonist of NPY Y5 receptors.

Table 1. Activity of hydrazide derivatives at the Y1, Y2, and Y5 NPY receptors.


Compd.	R ₁	R ₂	Biological Activity			
			% Inhib. Y ₁	% Inhib. Y ₂ ^b	% Inhib. Y ₂ ^c	% Inhib. Y ₅
1	-H	Ph-NH-	A>5·10 ⁻⁶ M	A>5 10 ⁻⁶ M	I	A>5·10 ⁻⁶ M
2	-H	Ph-CO-NH-	I	A>5·10 ⁻⁶ M	I	A>5·10 ⁻⁶ M
3	-H	2-Indolyl-CO-NH-	I		I	I
4	-H	3-Pyridyl-CO-NH-	I	I		24% 10 ⁻⁵ M
5	-H	3-Indolyl-CH ₂ -CO-NH-	10% 10 ⁻⁵ M			36% 10 ⁻⁵ M
6	-H	3-Pyrazolyl-	I	I		I
7	-OCH ₃	Ph-NH-	A>5·10 ⁻⁶ M	A>5 10 ⁻⁶ M	I	I
8	-OCH ₃	Ph-CO-NH-	A>5·10 ⁻⁶ M	I	I	I
9	-OCH ₃	2-Indolyl-CO-NH-	I	>10 ⁻⁵	I	I
10	-OCH ₃	3-Pyridyl-CO-NH-	I	I		22% 10 ⁻⁵ M
11	-OCH ₃	Ph-SO ₂ -NH-	23% 10 ⁻⁵ M			I
12	-OCH ₃	3-Indolyl-CH ₂ -CO-NH-	I			I
13	-OCH ₃	3-Pyrazolyl-	I			I

I= inactive
 A= agonist
 Y₁= rabbit saphenous vein
 Y₂^b= dog saphenous vein
 Y₂^c= rat deferens vessel
 Y₅= rabbit ileum

REFERENCES

- Gehlert, D. R.; *Proc. Soc. Exp. Biol. Med.* **1998**, 218, 7.
- Blomqvist, A. G.; Herzog, H.; *Trends Neurol Sci.* **1997**, 20, 294.
- Sahu, A.; Sninsky, C. A.; Kalra, Sp.; *Brain Res.* **1997**, 755, 339.
- Zimanyi, I. A.; Fathi, Z.; Poindexter, G. S.; *Curr. Pharm. Des.* **1998**, 4, 349.
- Kanataki, A.; Ishihara, A.; Asahi, S.; Tanaka, T.; Ozaki, S.; Ihara, M.; *Endocrinology* **1996**, 137, 3177.
- Doods, H. N.; Wieland, H. A.; Engel, W.; *Regul. Pept.* **1996**, 65, 71.
- Leoluca, C. *et al.*; *J. Clin. Invest.* **1998**, 102, 2136.
- Rueger, H.; Schmidlin, T.; Rigollier, P.; Yamaguchi, Y.; **1997**, WO97/20283-A2.
- Hong Pheng, L.; Quirion, R.; Iyengar, S.; Fournier, A.; Regoli, D.; *Eur. J. Pharmacol.* **1997**, 333, R3-R5.
- Serradeil-Le Gal, C.; Valette, G.; Rouby, P. E.; Pellet, A.; Oury-Donat, F.; Brossard, G.; Lespy, L.; Marty, E.; Neliat, G.; de Cointet, P.; Maffrand, J. P.; Le Fur, G.; *FEBS Lett.* **1995**, 362, 192.
- Fuhlendorff, J.; Gether, U.; Aakelund, L.; Langeland-Johansen, N.; Thogersen, H.; Melberg, S.G.; Olsen, U.B.; Thatrup, O.; Schwartz, T. W.; *Proc. Natl. Acad. Sci.* **1990**, 87, 182.
- Hu, Y.; Bloomquist, B. T.; Cornfield, L. J.; DeCarr, L. B.; Flores-Riveros, J.; Friedman, L.; Jiang, P.; Lewis-Higgins, L.; Sadlowski, Y.; Schaefer, J.; Velazquez, N.; McCaleb, M. L.; *J. Biol. Chem.* **1996**, 271, 26315.
- Mizayawa, T.; Otomatsu, T.; Yamada, T.; Kuwata, S.; *Tetrahedron Lett.* **1984**, 25, 771.
- Feletou, M. *et al.*; *Regulatory Peptides* **1998**, 75-76, 221